

# A Study of Burst Abdomen: It's Causes and Management

Dr. Brijesh Parekh<sup>1</sup>, Dr. Chetan Sharma<sup>2</sup>, Dr. Pooja Shah<sup>3</sup>, Dr. Prayas Bachani<sup>4</sup>

<sup>1</sup>MS, Associate Professor, Department of General Surgery, Sheth L. G. Hospital, Affiliated to AMCMET Medical College, Maninagar, Ahmedabad, Gujarat, India

<sup>2, 3, 4</sup>Resident doctors in department of General Surgery, Sheth L. G. Hospital, Affiliated to AMCMET Medical College, Maninagar, Ahmedabad, Gujarat, India

**Abstract:** ***Background:** Burst abdomen (abdominal wound dehiscence) is a severe post-operative complication. Burst abdomen is defined as post-operative separation of abdominal musculo-aponeurotic layers. The study aims to find etiological factors of burst abdomen in hospitalised patients, evaluate current management methods and to compare conservative and operative approach with respect to complication and outcomes. **Methods:** All cases presenting with abdominal wound dehiscence after surgery were included. An elaborate clinical history was taken in view of the significant risk factors, the types of surgery performed, type of disease involved and management methods and their outcome. A total of 30 cases were included in this prospective study. Data was analyzed using appropriate software. **Results:** The results concluded that male patients have a higher incidence of laparotomy wound dehiscence and in 5<sup>th</sup> decade. Patients presenting with peritonitis secondary to gastro-duodenal perforation are more prone to burst abdomen. **Conclusions:** Burst abdomen is a serious sequel of impaired wound healing. Presence of anaemia, hypoproteinaemia favours high incidence of burst abdomen. Delayed suturing, of burst abdomen has a lower frequency of complications. Adherence to proper technique and sincere efforts to minimize the impact of the predisposing factors play a much larger role in both treatment and prevention of this condition.*

**Keywords:** Abdominal wound dehiscence, Burst abdomen, Laparotomy

## 1.Introduction

Burst abdomen (abdominal wound dehiscence) is a severe post-operative complication. Incidence as described in literature ranges from 0.4% to 3.5%<sup>(1)</sup>. Burst abdomen is defined as post-operative separation of abdominal musculo-aponeurotic layers, which is recognised within days after surgery and requires some form of intervention. Various risk factors are responsible for wound dehiscence such as emergency surgery, intra-abdominal infection, malnutrition (hypoalbuminemia, anaemia), advanced age, systemic diseases (uraemia, diabetes mellitus) etc<sup>2</sup>. Good knowledge of these risk factors is mandatory for prophylaxis<sup>3</sup>. Patient identified as being high risk may benefit from close observation and early intervention. The study aims to find etiological factors of burst abdomen in hospitalised patients, evaluate current management methods and to compare conservative and operative approach with respect to complication and outcomes.

## 2.Methods

This is a prospective study carried out from December 2020 to December 2022 in the Department of General Surgery, Narendra Modi Medical College, Maninagar, Ahmedabad, Gujarat, India.


Total 30 patients who underwent both emergency or elective abdominal procedure and developed post-operative dehiscence during the study period were included. The inclusion criteria used were patients above 18 years of age of either sex, who gave consent for investigations and treatment. Exclusion criteria being primarily operated outside or patient who had undergone previous laparotomy for any condition (or had an incisional hernia or burst abdomen).

A comprehensive history and thorough physical examination with any other relevant history were recorded. Statistical analysis was processed using Excel software programs. Observations are represented as bar diagrams and pie charts.

## 3.Results


### Age

The youngest patient was 21 years old and the oldest patient was 70 years old. The highest incidence of burst abdomen in the present study was between 51 and 60 years of age, the average age being approximately 49 years. The patients in this study were in the range of 49±13.5 (standard deviation) years.



**Figure 1:** Age and number of study participants with burst abdomen

**Sex distribution**




**Figure 2:** Sex Distribution in Cases of Burst Abdomen


In present study, 25 patients (83.3%) of the patients were male and the remaining 5 (16.7 %) were females. The male: female ratio was approximately 5: 1.

**Preoperative predisposing causes**

The study showed that the majority of patients had intra-abdominal sepsis (24 patients) and anaemia (22 patients) as preoperative predisposing factors. Many patients had more than one predisposing factor.



**Figure 3:** Incidence of Burst Abdomen in Planned and Emergency Cases



**Figure 4:** Post-Operative Wound Discharge

**Planned or emergency surgery**

The incidence of burst abdomen was much higher in patients operated as emergency surgery (27/30) as compared to planned surgery (3/30).

**Intra-abdominal pathology and its origin**

Indication of laparotomy being perforation peritonitis are most commonly being gastro duodenal perforation (29.26%) and Ileal perforation (19.51%) other indication.

**Type of closure**

Mass closure was the standard technique used in all the cases in the series; the technique involves incorporating all of the layers of the abdominal wall (except skin) as one structure. Continuous sutures with No.1 Polyamide were used in 20 patients, in other 10 patients; abdomen was closed with simple interrupted Polyamide sutures.

**Time of disruption**

The majority of burst abdomen occurred between 7<sup>th</sup> and 10<sup>th</sup> post-operative day, with the highest incidence on the 7<sup>th</sup> post-operative day.

**Post-operative wound discharge**

In present study, 24 patients out of 30 had serosanguinous discharge from the wound.4 patients out of 30 had purulent discharge.2 patients experienced feculent discharge from wound site.

**Partial or complete burst**

In present study, 18 patients out of 30 (60%) had complete burst involving the whole length of the wound while 12 patients out of 30 (40%) had partial burst.

**Culture of discharge**

Gram-negative organisms were the ones most commonly grown from culture of the wound discharge.

**Management**

Conservative treatment (daily saline dressings) was done in 4/30 cases. Immediate resuturing of the wound in the operation theatre was done in 5 patients with tension sutures. Through and through vertical mattress sutures with supporting PVC tubing were used.

While 7 patients were re sutured without tension sutures. In 14 patient delayed secondary suturing was done.

**Complications of management and their treatment**

Out of 30 patients 14 (46.7%) patients fully recovered, 8 patient developed incisional hernia over a period of 3 to 21 months, 3 patients developed subsequent re-burst who subsequently died due to septicaemia. Out of 30 patients in the present study death occurred in 8 patients (27%) of

which 1 patients was from group I, 2 patient was from group II, 3 patients were from group III while 2 patients were from group IV.

### Total hospital stay

Total hospital stay of the patients increases because of burst abdomen. In present study, out of 30 patients, 5 had a total hospital stay between 31 and 40 days. The mean duration of total hospital stay was  $35 \pm 6.9$  days.

### Mortality

Out of 30 patients, death occurred in 8 patients, giving a mortality rate of 27%. All 8 died of complications due to septicaemia and multi-organ failure.

## 4. Discussion

This study reviewed 30 patients who had laparotomy wound dehiscence over a period from December 2020 to December 2022. Present study analyzed the potential causes of burst abdomen.

In present study average age of burst abdomen is 49 years. Burst abdomen is more common in males. In this study 83% cases were male which is well comparable with other studies.

High incidence of burst abdomen was seen in emergency operation as compared to elective surgeries. Lack of bowel preparation, preoperative optimization and higher frequency of contaminated cases are the major causes responsible for burst abdomen in emergency surgeries. In present study 27 cases underwent emergency operation while 3 cases underwent elective cases.

Present study showed that peritonitis due to perforation was a common cause of burst abdomen. Amongst which gastro duodenal perforation accounted for 29.26%. Other intra-abdominal pathologies in present study are ileal perforation (19.51%), Intestinal obstruction (18.29%), malignancy (14.63%), Koch's abdomen (2.43%), Stab injury (4.87%) and blunt trauma abdomen (3.65%). Peritonitis was the most common cause associated with burst abdomen.

In 22 patients out of 30 cases haemoglobin level was less than 10 gm%. The incidence of anaemia in cases of burst abdomen varies widely from series to series. It was only 6.66% in the study conducted by Wolf WI et al, while it was 90% in study conducted by Pierre J et al and 100% in study conducted by Waqar SH et al. (5, 8, 9). Hypoproteinaemia is also one of the most important factor which leads to delayed wound healing. In present study 18 patients out of 30 cases had a serum protein below 6 gm%. Afzal et al studies a subgroup of patients with peritonitis had protein energy malnutrition as one of significant risk factor for burst abdomen.

Out of 30 patients 8 developed incisional hernia. Many surgeons place retention sutures at laparotomy closure, in those with several risk factors for burst abdomen. Despite these measures, repaired dehisced laparotomy wound have a 69% incisional hernia development risk (over 10 years period), the majority of which develops over first 2 years.

There was a 27% mortality in present study. Although the incidence of burst abdomen has not changed much, the mortality due to it has decreased due to early recognition, early ambulation, better broad-spectrum antibiotics, better post-operative management and increasing awareness about the condition.

**Table 1:** Mode of Given to the Patients of Burst Abdomen

Group	Treatment given	Percentage	No. of cases
I	Immediate resuturing with tension suture	15.49%	5
II	Immediate resuturing without tension suture	21.95%	7
III	Delayed secondary suturing	47.56%	14
IV	Conservative management	14.63%	4

**Table 2:** Predisposing Factors Observed in Present Study, Out of 30 Cases

Predisposing factors	No. of cases
Intra abdominal sepsis	24
Anemia	22
Hypoproteinemia	18
Chest disease	16
Diabetes	9
Uraemia	10
Jaundice	6

**Table 3:** Frequency of Pathologies among Patients

Indication	Percentage	Cases
Gastro duodenal perforation	29.26%	9
Ileal perforation	19.51%	6
Intestinal obstruction	18.29%	5
Malignancy	14.63%	4
Large bowel perforation	7.31%	2
Koch's abdomen	2.43%	1
Stab injury	4.87%	1
Blunt trauma abdomen	3.65%	2

**Table 4:** Comparison of Outcome in the Four Management Groups

	GROUP I	GROUP II	GROUP III	GROUP IV	PERCENTAGE	No. of cases
Full recovery	4	2	8	0	46.67%	14
Incisional hernia	1	0	4	3	26.67%	8
Re-burst	0	0	3	0	10%	3
Death	1	2	3	2	26.67%	8

## 5. Conclusions

Burst abdomen is a serious sequel of impaired wound healing. Presence of anaemia, hypoproteinaemia favours high incidence of burst abdomen. Gram negative bacteria are most common organism involved in abdominal wound dehiscence. When operative and conservative treatment was compared, it was found that retention suture placement helped to decrease the frequency of complications when immediate resuturing was performed. The conservative approach had a higher morbidity. Hence delayed suturing, which had a lower frequency of complications in this study, may serve as a "middle path" between the two options. Burst abdomen remains a dreaded post-operative complication. Newer materials and devices continue to be developed and may simplify the treatment of burst abdomen, but adherence to proper technique and sincere efforts to minimize the impact of the predisposing factors play a much larger role in both treatment and prevention of this condition.

## References

- [1] Afzal S, Bashir MM. Determinants of wound dehiscence in abdominal surgery in public sector hospital. Annals of King Edward Medical University. 2010 Apr 1; 14 (3): 119.
- [2] Robert J, Fittzgebons JR. Nyhus and Condons hernia. Diagnostic and Imaging of abdominal wall hernia, 5<sup>th</sup> Ed., Lippincott Williams; 2002.
- [3] Gabrielle H, van Ramshorst, Nieuwenhuizen J, Hop WCJ, Arends P, Boom J, et al. Abdominal wound dehiscence in adults: development and validation of a risk model. World J Surg. 2010; 34: 20-7.
- [4] Spiliotis J, Konstantino S, Siveriotis T, Datsis AD, Archodaula, Georgios, et al. Wound dehiscence. World J Emerg Surg. 2009; 4: 12.
- [5] Waqer S, Malik Z, Razzaq A, Abdullah MT, Shaima A, Zahid MA. Frequency and risk factors for wound dehiscence/burst abdomen in midline laparotomies. Journal Ayub Med Coll. 2005; 17 (4): 70-3.
- [6] Lotfy W. Burst Abdomen: Is it a preventable complication?. Egyptian J Surg. 2009; 28 (3): 128-32.
- [7] Halasz NA. Dehiscence of laparotomy wounds. Am J Surg. 1968; 116 (2): 210-4.
- [8] Riou JP, Cohen JR, Johnson H. Factors influencing wound dehiscence. Am J Surg. 1992; 163 (3): 324-30.
- [9] Wolff WI. Disruption of abdominal wounds. Ann of Surg. 1950; 131: 534-55
- [10] Afzal S, Bashir MM. Determinants of wound dehiscence in abdominal surgery in public sector hospital. Ann King Edward Med University. [11] 2010; 14 (3): 119.
- [12] Hartzell JB, Winfield JM, IRVIN JL. Plasma Vitamin C and serum protein levels in wound disruption. J Am Med Association 1941; 116 (8): 669-74