

A Discussion on the Novel Concept of b^+ Open Sets in Simple Extended Topological Spaces

R. Sujeetha

PG Student, Department of Mathematics, Nirmala College for Women, Coimbatore, India

Email: [sujeearg0320\[at\]gmail.com](mailto:sujeearg0320[at]gmail.com)

Abstract: *The purpose of this article is to introduce, and study the newly proposed concept of B^+C open set in simple extension topology. In this article we discuss some of the properties of B^+C open set, and obtain certain characterization and preserving theorems of B^+C -interior and B^+C -closure.*

Keywords: b^+ open set, B^+C -open set

1. Introduction

The class of generalized open sets in a topological space is called b -open sets was introduced by Andrijevic [1]. H. Z. Ibrahim [2] introduced the concept of a BC -open set. In 1968, Velicko [10] introduced the concept of θ -open. Di Maio and Noiri [11] introduced the concept of semi- θ -open. R. H. Yunis [12] introduced the concept of properties of θ -semi open sets. In 1963, Levine [5] introduced the concept of simple extension of a topology $\tau(B) = \{(B \cap O) \cup O' / O, O' \in \tau\}$. F. Nirmala Irudayam [3] introduced the concept of b^+ -open sets in extended topological spaces. The class of b^+ -open sets is contained in the class of semi- pre^+ open sets and contains all semi- pre^+ open sets. Joseph and Kwack [9] introduced the concept of θ -semi open sets using semi-open sets. It is well-known that a space X is called T_1 if for, each pair of distinct points x, y of X there exists a pair of open sets, one containing x but not y and the other containing y but not x , as well as is T_1 if and only if for any point $x \in X$, the singleton set $\{x\}$ is closed. In this article, (X, τ) stand for topological spaces with no separation axioms assumed unless otherwise stated.

2. Preliminaries

Definition 2.1

A subset A of a space X is called b^+ open if $A \subseteq \text{Int}(\text{Cl}^+(A) \cup \text{Cl}^+(\text{Int}(A)))$. The family of all b^+ open subsets of a simple extended topological space (X, τ^+) is denoted by $B^+O(X, \tau^+)$ or (Briefly, $B^+O(X)$).

Lemma: 2.2

For a subset A of a space (X, τ) , the following conditions are equivalent:

- 1) $A \in RO(X)$.
- 2) $A \in \tau \cap SC(X)$.
- 3) $A \in \alpha O(X) \cap SC(X)$.
- 4) $A \in PO(X) \cap SC(X)$.

Definition: 2.3

A subset A of a space X is called θ -semi-open if for each $x \in A$, there exist a semi-open set G such that each $x \in G \subset \text{Cl}(G) \subset A$.

Definition: 2.4

A subset A of a space X is called **semi- θ -open** if for each $x \in A$, there exist a semi-open set G such that each $x \in G \subset S\text{Cl}(G) \subset A$.

Definition: 2.5

A subset A of a space X is called **θ -open** if for each $x \in A$, there exist an open set G such that each $x \in G \subset \text{Cl}(G) \subset A$.

Theorem: 2.6

If X is s^{**} -normal, then $S\theta O(X) = \theta O(X) = \theta SO(X)$.

We recall that a topological space X is said to be extremely disconnected, if $\text{Cl}(G)$ is open for every open set G of X .

Definition: 2.7

A space X is called locally indiscrete if every open subset of X is closed.

Theorem: 2.8

A space X is extremely disconnected if and only if $\delta O(X) = \theta SO(X)$.

Theorem: 2.9

A space X is extremely disconnected if and only if $RO(X) = RC(X)$.

3. On B^+C -Open set

This section introduces a new class of b^+ -open sets namely B^+C -open sets in simple extended topological spaces and various properties of this novel set are studied.

Definition: 3.1

A subset A of a space (X, τ^+) is called B^+C open, if for each $x \in A \in B^+O(X)$, there exists a closed set F such that $x \in F \subset A$.

The family of all B^+C open subsets is denoted by $B^+CO(X)$ of a simple extended topological space (X, τ^+) .

Theorem: 3.2

A subset A of a space X is B^+C open, if and only if A is b^+ open and it is a union of closed sets in (X, τ^+) . i.e. $A = \cup F_\alpha$.

That is $A = \bigcup F_\alpha$ where A is b^+ open set, and F_α is closed sets in (X, τ^+) for each α .

Proof:

The proof is obvious from the definition.

Remark: 3.3

Every B^+C open subset of a space (X, τ^+) is b^+ open.

Remark: 3.4

The converse of the above remark need not be true, as shown in the following example,

Example:3.5

Consider $X = \{1, 2, 3\}$ with the topology

$$\tau = \{\varphi, X, \{1\}, \{2\}, \{1,2\}\}$$

$$\tau^c = \{\varphi, X, \{2,3\}, \{1,3\}, \{3\}\}$$

$$B = \{3\}, \tau^+(B) = \{(B \cap O) \cup O / O, O \in \tau\}$$

Then the family of closed sets are:

$$\{\varphi, X, \{1\}, \{2\}, \{2,3\}, \{1,3\}, \{2,3\}\}$$

Hence from the definitions we find the following families:

$$B^+O(X) = \{\varphi, X, \{1\}, \{2\}, \{1,2\}, \{1,3\}, \{2,3\}\} \text{ and}$$

$$B^+CO(X) = \{\varphi, X, \{1,3\}, \{2,3\}\}$$

Theorem:3.6

Let $\{A_\alpha : \alpha \in \Delta\}$ be a collection of B^+C open sets in a simple extended topological space (X, τ^+) , then $\bigcup \{A_\alpha : \alpha \in \Delta\}$ is B^+C open.

Proof:

Let A_α be a B^+C open set for each α , then A_α is b^+ open.

Hence $\bigcup \{A_\alpha : \alpha \in \Delta\}$ is b^+ open.

Let $x \in \bigcup \{A_\alpha : \alpha \in \Delta\}$, there exist $\alpha \in \Delta$ such that $x \in A_\alpha$.

Since A_α is b^+ open for each α , there exists a closed set F .

Such that $x \in F \subset A_\alpha \subset \bigcup \{A_\alpha : \alpha \in \Delta\}$.

So $x \in F \subset \bigcup \{A_\alpha : \alpha \in \Delta\}$.

Therefore, $\bigcup \{A_\alpha : \alpha \in \Delta\}$ is B^+C open set.

Theorem:3.7

If the family of all b^+ open sets of a space X is a topology on X , then the family of B^+C open sets is also a topology on X .

Proof:

Clearly $\varphi, X \in B^+CO(X)$ and by Theorem 3.6 the union of any family of B^+C open sets is B^+C open.

To complete the proof it is enough to show that the finite intersection of B^+C open sets is B^+C open set.

Let A and B be two B^+C open sets then A and B are b^+ open sets.

Since $B^+O(X)$ is a topology on X , so $A \cap B$ is b^+ open.

Let $x \in A \cap B$, then $x \in A$ and $x \in B$, so there exists F and E such that $x \in F \subset A$ and $x \in E \subset B$ this implies that $x \in F \cap E \subset A \cap B$.

Since intersection of closed sets is closed, $F \cap E$ is closed set.

Thus $A \cap B$ is B^+C open set.

This completes the proof.

Theorem:3.8

The set A is B^+C open in the simple extended topological space (X, τ^+) , if and only if for each $x \in A$, there exists a B^+C open set B such that $x \in B \subset A$.

Proof:

Assume that A is B^+C open set.

Then for each $x \in A$, put $A = B$ is B^+C open set containing x such that $x \in B \subset A$.

Conversely, suppose that for each $x \in A$, there exists a B^+C open set B such that $x \in B \subset A$

Thus $A = \bigcup B_x$ where $B_x \in B^+CO(X)$ for each x , therefore A is B^+C open set

In the following theorem, the family of b^+ -open sets is identical to the family of B^+C open sets.

Theorem:3.9

If a space (X, τ^+) is T_1 -space, then the families $B^+O(X) = B^+CO(X)$.

Proof:

Let A be any subset of a space (X, τ^+) and $A \in B^+O(X)$.

If $A = \varphi$, then $A \in B^+CO(X)$.

If $A \neq \varphi$, then for each $x \in A$.

Since space X is T_1 , then every singleton in (X, τ^+) is closed set.

Hence $x \in \{x\} \subset A$ and $\{x\} \in B^+CO(X)$.

Thus $B^+O(X) \in B^+CO(X)$, but $B^+CO(X) \subset B^+O(X)$

Therefore $B^+O(X) = B^+CO(X)$.

Definition: 3.10

1) Let A be a subset of a simple extension topological space (X, τ^+) is called θ^+ - open set if for each $x \in X$, there exists an open set G such that $x \in G \subset (Cl^+(G)) \subset A$.

2) Let A be a subset of a simple extension topological space (X, τ^+) is said to be θ^+ - semi open set if there exists a θ^+ - open set U of X such that $U \subset A \subset Cl^+(U)$.

3) Let A be a subset of a simple extension topological space (X, τ^+) called δ^+ - open set, if for each $x \in A$, there exists an open set G such that $x \in G \subset Int(Cl^+(G)) \subset A$.

4) In simple extension topological space (X, τ^+) to be s^{**} -normal if and only if for every semi $^+$ closed set F and every semi $^+$ open set G containing F , there exist an open set H such that $F \subset H \subset Cl^+(H) \subset G$.

5) Let A be a subset of a topological space (X, τ^+) is **Regular $^+$ -Open** if $A = Int(Cl^+(A))$.

6) Let A be a subset of a topological space (X, τ^+) is **Regular $^+$ -Closed** if $A = Cl^+(Int(A))$.

Theorem:3.11

Every θ^+ -semi open set of a space (X, τ^+) is B^+C open set.

Proof:

Let A be a θ^+ -semi open set in (X, τ^+) , then for each $x \in A$, there exists a semiopen set G . Such that $x \in G \subset Cl^+(G) \subset A$, so $\bigcup \{x\} \in \bigcup G \cup Cl^+(G) \subset A$ for each $x \in A$ implies that $A = \bigcup Cl^+(G)$, which is semi-open set and $A = \bigcup Cl^+(G)$ is a union of closed sets, by Theorem 3.2, A is B^+C open set.

The following corollaries are the direct implications from the definition 3.10-(5) and (6).

Corollary:3.12

- 1) Every θ^+ -open set is B^+C open.
- 2) Every regular $^+$ -closure is B^+C open set.

Theorem:3.13

If a simple extended topological space (X, τ^+) is locally indiscrete, then $S^+O(X) \subset B^+CO(X)$.

Proof:

Let A be any subset of a space X
 $A \in S^+O(X)$, if $A = \emptyset$ then $A \in B^+CO(X)$,
 If $A \neq \emptyset$, then $A \subset Cl^+Int(A)$.
 Since X is locally indiscrete, then $Int(A)$ is closed
 Hence $Int(A) \subset A$, this implies that for
 each $x \in A$, $x \in (Int(A) \subset A)$.
 Therefore, A is B^+C open set.
 Hence $S^+O(X) \subset B^+CO(X)$.

Remark: 3.14

Every open set in (X, τ^+) is semi $^+$ -open, it follows that if a simple extended topological space (X, τ^+) is T_1 or locally indiscrete, then $\tau^+ \subset B^+CO(X)$.

Theorem: 3.15

Let (X, τ^+) be a simple extended topological space, if X is regular, then $\tau^+ \subset B^+CO(X)$.

Proof:

Let A be any subset of a simple extended topological space (X, τ^+) .
 Consider A be an open, if $A = \emptyset$, then $A \in B^+CO(X)$.
 If $A \neq \emptyset$, since X is regular, so for each $x \in A \subset X$, there exists an open set G such that
 $x \in G \subset Cl^+(G) \subset A$.
 Thus we have $x \in Cl^+(G) \subset A$.
 Since $A \in \tau^+$ and hence $A \in B^+O(X)$,
 Therefore $\tau^+ \subset B^+CO(X)$.

Theorem 3.16

Let (X, τ^+) be an extremely disconnected space and if $A \in \delta^+O(X)$, then $A \in B^+CO(X)$.

Proof:

Let $A \in \delta^+O(X)$.
 If $A = \emptyset$, then $A \in B^+CO(X)$.
 If $A \neq \emptyset$, since a space X is extremely disconnected.
 Then $\delta^+O(X) = \theta^+SO(X)$
 Hence $A \in \theta^+SO(X)$.
 But $\theta^+SO(X) \subset B^+CO(X)$
 Therefore, $A \in B^+CO(X)$

Theorem: 3.17

Let (X, τ^+) be an extremely disconnected space, if $A \in R^+O(X)$, then $A \in B^+CO(X)$.

Proof:

The above theorem can be proved easily using theorem 3.26, and the conditional that

$$R^+O(X) \subset \delta^+O(X)$$

Theorem:3.18

Let (X, τ^+) be an s^{+**} -normal space. If $A \in S^+O(X)$, then $A \in B^+CO(X)$.

Proof:

Let $A \in S^+O(X)$.
 If $A = \emptyset$, then $A \in B^+CO(X)$.
 If $A \neq \emptyset$, since space X is s^{+**} -normal, $S^+O(X) = \theta^+SO(X)$.
 Hence $A \in \theta^+SO(X)$.
 But $\theta^+SO(X) \in B^+CO(X)$.
 Therefore, $A \in B^+CO(X)$.

Theorem: 3.19

For any subset A of a simple extended topological space (X, τ^+) and $B^+O(X) = S^+\theta(X)$.

The following conditions are equivalent:

- i) A is regular $^+$ closed.
- ii) A is closed and B^+ Copen.
- iii) A is closed and b^+ open.
- iv) A is α -closed and b^+ open.
- v) A is pre-closed and b^+ open.

Definition:3.20

A subset B of a space X is called B^+C closed, if X/B is B^+C open. The family of all B^+C -closed subsets of a simple extended topological space (X, τ^+) is denoted by $B^+CC(X, \tau^+)$ or (Briefly, $B^+CC(X)$).

Theorem:3.21

A subset B of a space X is B^+C -closed, if and only if B is a b^+ -closed set, and it is an intersection of open sets.

Proof: Proof is straight forward.

Theorem:3.22

Let $\{b^+_{\alpha} : \alpha \in \Delta\}$ be a collection of B^+C -closed sets in a topological space (X, τ^+) .
 Then $\cap \{b^+_{\alpha} : \alpha \in \Delta\}$ is B^+C -closed.

Proof: The proof is analogous to theorem 3.6

The union of two B^+C -closed sets need not be B^+C closed as is shown by the following counterexample,

Example: 3.23

In Example 3.5, the family of B^+C closed subset of X is $B^+CC(X) = \{\emptyset, X, \{1\}, \{2\}\}$. Here $\{A\} \in B^+CC(X)$ and $\{B\} \in B^+CC(X)$, but $\{1\} \cup \{2\} = \{1,2\} \notin B^+CC(X)$.
 All of the following results are true by using complement.

Result: 3.24

If a space (X, τ^+) is T_1 , then $B^+CC(X) = B^+C(X)$.

Result: 3.25

For any subset B of a space (X, τ^+) . If $B \in \theta^+SC(X)$, then $B \in B^+CC(X)$.

Corollary: 3.26

Each θ^+ -closed set is B^+C closed.

Corollary: 3.27

Each regular⁺open set is B^+C closed.

Remark: 3.28

If a simple extended topological space (X, τ^+) is locally indiscrete, then $S^+C(X) \subset B^+CC(X)$.

Corollary: 3.29

Let (X, τ^+) be a simple extended topological space, if X is regular or locally indiscrete, then the family of closed sets is a subset of the family of B^+C -closed sets.

Corollary: 3.30

Let (X, τ^+) be any extremely disconnected space. If $B \in \delta^+C(X)$, then $B \in B^+CC(X)$.

Corollary: 3.31

Let (X, τ^+) be an extremely disconnected space. If $B \in R^+C(X)$, then $B \in B^+CC(X)$.

Corollary: 3.32

Let (X, τ^+) be a s^{**} -normal space. If $B \in S^+\theta C(X)$, then $B \in B^+CC(X)$.

Theorem: 3.33

For any subset B of a space (X, τ^+) and $S^+C(X) = B^+C(X)$.

The following conditions are equivalent:

- i) b^+ is regular⁺open.
- ii) b^+ is open and B^+C -closed.
- iii) b^+ is open and b^+ -closed.
- iv) b^+ is α -open and b^+ -closed.
- v) b^+ is preopen and b^+ -closed.

4. Some Properties of B^+C Open Sets

This section is devoted to the study of B^+C -neighborhood, B^+C -interior, B^+C -closure of B^+C derived set via the newly coined B^+C -open sets.

Definition: 4.1

Let (X, τ^+) be a simple extended topological space and $x \in X$, then a subset N of x is said to be a b^+C -neighborhood of x , if there exists a B^+C -open set U in X such that $x \in U \subset N$.

Theorem: 4.2

In a simple extended topological space (X, τ^+) , a subset A of X is B^+C -open, if and only if it is a B^+C -neighbourhood of each of its points.

Proof:

Let $A \subset X$ be a B^+C -open set, since for every $x \in A$, $x \in A \subset A$ and A is B^+C -open, this shows that A is a B^+C -neighbourhood of each of its points.

Conversely, suppose that A is a B^+C -neighbourhood of each of its points.

Then for each $x \in A$, there exists $b^+x \in B^+CO(X)$ such that $b^+x \subset A$.

Then $A = \cup \{b^+x : x \in A\}$.

Since each b^+x is B^+C -open.

It follows that A is B^+C -open set.

Theorem: 4.3

For any two subsets A, B of a simple extended topological space (X, τ^+) and $A \subset B$, if A is a B^+C -neighbourhood of a point $x \in X$, then B is also B^+C -neighbourhood of the same point x .

Proof:

Let A be a B^+C -neighbourhood of $x \in X$ and $A \subset B$ then by Definition 3.1.

There exists a B^+C -open set U such that $x \in U \subset A \subset B$.

This implies that B is also a B^+C -neighbourhood of x .

Remark: 4.4

Every B^+C neighborhood of a point is ab^+ -neighbourhood, This follows from the fact that every B^+C -open set is b^+ -open.

Definition: 4.5

Let A be a subset of a topological space (X, τ^+) , a point $x \in X$ is said to be B^+C -interior point of A , if there exist a B^+C -open set U such that $x \in U \subset A$. The set of all B^+C -interior points of A is called B^+C -interior of A and is denoted by $B^+C \text{Int}(A)$.

Some properties of the B^+C -interior of a set are investigated in the following theorem.

Theorem: 4.6

For subsets A, B of a space X , the following statements hold,

- i) $B^+C \text{Int}(A)$ is the union of all B^+C -open sets which are contained in A .
- ii) $B^+C \text{Int}(A)$ is B^+C -open set in X .
- iii) A is B^+C -open if and only if $A = B^+C \text{Int}(A)$.
- iv) $B^+C \text{Int}(B^+C \text{Int}(A)) = B^+C \text{Int}(A)$.
- v) $B^+C \text{Int}(\phi) = \phi$ and $B^+C \text{Int}(X) = X$.
- vi) $B^+C \text{Int}(A) \subset A$.
- vii) If $A \subset B$, then $B^+C \text{Int}(A) \subset B^+C \text{Int}(B)$.
- viii) If $A \cap B = \phi$, then $B^+C \text{Int}(A) \subset B^+C \text{Int}(B)$.
- ix) $B^+C \text{Int}(A) \cup B^+C \text{Int}(B) \subset B^+C \text{Int}(A \cup B)$.
- x) $B^+C \text{Int}(A \cap B) \subset B^+C \text{Int}(A) \cap B^+C \text{Int}(B)$.

Proof:

Let $x \in X$ and $x \in B^+C \text{Int}(A)$, then by Definition 4.5

There exists a B^+C -open set U such that $x \in U \subset A \subset B$ implies that $x \in U \subset B$.

Thus $x \in B^+C \text{Int}(B)$.

The other parts of the theorem can be proved easily.

Theorem: 4.7.

For a subset A of a simple extended topological space (X, τ^+) , then

$B^+C \text{Int}(A) \subset b^+ \text{Int}(A)$.

Proof:

This follows immediately since all B^+C -open set is b^+ -open.

Definition: 4.8.

Let A be a subset of a space X . A point $x \in X$ is said to be B^+C -limit point of A , if for each B^+C open set U containing x , $U \cap (A \setminus \{x\}) \neq \phi$. Then the set of all B^+C -limit points of

A is called a B^+C -derived set of A, and is denoted by $B^+CD(A)$.

Theorem:4.9.

Let A be a subset of X, if for each closed set F of X containing x such that

$F \cap (A \setminus \{x\}) \neq \varnothing$, then a point $x \in X$ is B^+C -limit point of A.

Proof:

Let U be any B^+C -open set containing x.

Then for each $x \in U \in B^+O(X)$, there exists a closed set F such that $x \in F \subset U$.

By hypothesis, we have $F \cap (A \setminus \{x\}) \neq \varnothing$.

Hence $U \cap (A \setminus \{x\}) \neq \varnothing$.

Therefore, a point $x \in X$ is B^+C -limit point of A.

Some properties of B^+C -derived set are stated in the following theorem,

Theorem:4.10.

Let A and B be subsets of a space X. Then we have the following properties:

- i) $B^+CD(\varnothing) = \varnothing$
- ii) If $X \in B^+CD(A)$, then $x \in B^+CD(A \setminus \{X\})$.
- iii) If $A \subset B$, then $B^+CD(A) \subset B^+CD(B)$.
- iv) $B^+CD(A) \cup B^+CD(B) \subset B^+CD(A \cup B)$
- v) $B^+CD(A \cap B) \subset B^+CD(A) \cap B^+CD(B)$
- vi) $B^+CD(B^+CD(A)) \setminus A \subset B^+CD(A)$.
- vii) $B^+CD(A \cup B^+CD(A)) \subset (A \cup B^+CD(A))$

Proof: We only prove vi), vii), and the other results can be proved obviously.

vi) If $x \in B^+CD(B^+CD(A) \setminus A)$.

U is a B^+C -open set containing x.

$U \cap (B^+CD(A) \setminus \{x\}) \neq \varnothing$.

Let $y \in U \cap (B^+CD(A) \setminus \{x\})$

Since $y \in B^+CD(A)$ and $y \in U$, $U \cap (A \setminus \{y\}) \neq \varnothing$.

Let $z \in U \cap (A \setminus \{y\})$.

Then, $z \neq x$ for $Z \in A$ and $x \notin A$.

Hence $U \cap (A \setminus \{x\}) \neq \varnothing$.

Therefore, $x \in B^+CD(A)$.

vii) Let $x \in B^+CD(A \cup B^+CD(A))$. If $x \in A$ the result is obvious.

Let $x \in B^+CD(A \cup B^+CD(A) \setminus A)$.

Then for B^+C open set U containing x, $U \cap (A \cup B^+CD(A) \setminus \{x\}) \neq \varnothing$.

Thus, $U \cap (A \setminus \{x\}) \neq \varnothing$ or $U \cap (B^+CD(A) \setminus \{x\}) \neq \varnothing$.

Now, it follows similarly from i) that $U \cap (A \setminus \{x\}) \neq \varnothing$.

Hence, $x \in B^+CD(A)$.

Therefore, in any case $(B^+CD(A) \cup B^+CD(A)) \subset A \cup B^+CD(A)$.

Corollary:4.11.

For a subset A of a simple extended topological space (X, τ^+) , then $b^+D(A) \subset B^+CD(A)$

Proof:

It is sufficient to recall that every B^+C -open set is b^+ open.

Definition:4.12

For any subset A in a simple extended topological space (X, τ^+) , the B^+C -closure of A, denoted by $B^+CCl^+(A)$, is defined by the intersection of all B^+C -closed sets containing A.

Theorem:4.13

A subset A of a simple extended topological space (X, τ^+) is B^+C -closed if and only if it contains the set of its B^+C limit points.

Proof:

Assume that A is B^+C -closed

If possible that x is a B^+C -limit point of A which belongs to $X \setminus A$, then $X \setminus A$ is B^+C -open set containing the B^+C -limit point of A, therefore $A \cap X \setminus A \neq \varnothing$.

which is a contradiction.

conversely, assume that A contains the set of its B^+C limit points.

For each $x \in X \setminus A$, there exists a B^+C -open set U containing X such that $A \cap U = \varnothing$,

that is $x \in U \subset X \setminus A$ by Theorem 3.8, $X \setminus A$ is B^+C -open set

Hence A is B^+C -closed set.

Theorem:4.14.

Let A be a subset of a space (X, τ^+) , then $B^+CCl^+(A) = A \cup B^+CD(A)$.

Proof:

Since $B^+CD(A) \subset B^+CCl^+(A)$ and $A \subset B^+CCl^+(A)$.

Then $A \cup B^+CD(A) \subset B^+CCl^+(A)$, on the other hand,

to prove that $B^+CCl^+(A) \subset A \cup B^+CD(A)$.

since $B^+CCl^+(A)$ is the smallest B^+C -closed set containing A,

so it is enough to prove that $A \cup B^+CD(A)$ is B^+C -closed.

Let $x \notin A \cup B^+CD(A)$.

This implies that $x \notin A$ and $x \notin B^+CD(A)$.

Since $x \notin B^+CD(A)$, there exists a B^+C -open set $G(x)$ of x which contains no point of A other than x but $x \notin A$.

So $G(x)$ contains no point of A, which implies $G(x) \subset X \setminus A$.

Again, $G(x)$ is a B^+C -open set of each of its points.

But as $G(x)$ does not contain any point of A, no point of $G(x)$ can be a B^+C -limit point of A.

Therefore, no point of $G(x)$ can belong to $B^+CD(A)$.

This implies that $G(x) \subset X \setminus B^+CD(A)$.

Hence, it follows that $x \in G(x) \subset X \setminus A \cap X \setminus B^+CD(A) \subset X \setminus (A \cup B^+CD(A))$

Therefore, $A \cup B^+CD(A)$ is B^+C -closed.

Hence $B^+CCl^+(A) \subset A \cup B^+CD(A)$.

Thus $B^+CCl^+(A) = A \cup B^+CD(A)$.

Corollary:4.15.

Let A be a set in a space (X, τ^+) . A point $x \in U$ is in the B^+C -closure of A if and only if

$A \cap U \neq \varnothing$, for every B^+C -open set U containing x.

Proof:

Let $x \notin B^+CCl^+(A)$.

Then $x \notin F$, where F is B^+C -closed with $A \subset F$.

So $x \in X \setminus F$ and $X \setminus F$ is a B^+C -open set containing x .

Hence $(X \setminus F) \cap A \subset (X \setminus F) \cap (\cap F) = \emptyset$.

Conversely, suppose that there exists a B^+C -open set containing x with $A \cap U = \emptyset$.

Then $A \subset X \setminus U$ and $X \setminus U$ is a B^+C -closed.

Hence $x \notin B^+CCl^+(A)$.

Theorem:4.16

Let A be any subset of a space (X, τ^+) . If $A \cap U \neq \emptyset$ for every closed set F of X containing x , then the point x is in the B^+C -closure of A .

Proof:

Suppose that U be any B^+C -open set containing x , then by Definition 3.1, there exists a closed set F .

Such that $x \in F \subset U$, so by hypothesis, $A \cap F \neq \emptyset$ implies $A \cap U \neq \emptyset$ for every B^+C -open set U containing x .

Therefore $x \in B^+CCl^+(A)$.

Here we introduce some properties of B^+C -closure of the sets.

Theorem:4.17

For subsets A, B of a space (X, τ^+) , the following statements are true.

- 1) The B^+C -closure of A is the intersection of all B^+C closed sets containing A .
- 2) $A \subset B^+CCl^+(A)$.
- 3) $B^+CCl^+(A)$ is B^+C -closed set in X .
- 4) A is B^+C -closed set if and only if $A = B^+CCl^+(A)$.
- 5) $B^+CCl^+(B^+CCl^+(A)) = B^+CCl^+(A)$.
- 6) $B^+CCl^+(\emptyset) = \emptyset$ and $B^+CCl^+(X) = X$.
- 7) If $A \subset B$, then $B^+CCl^+(A) \subset B^+CCl^+(B)$.
- 8) If $B^+CCl^+(A) \cap B^+CCl^+(B) = \emptyset$, then $AB = \emptyset$.
- 9) $B^+CCl^+(A) \cup B^+CCl^+(B) \subset B^+CCl^+(A \cup B)$.
- 10) $B^+CCl^+(A \cap B) \subset B^+CCl^+(A) \cap B^+CCl^+(B)$.

Proof: Obvious.

Theorem:4.18

For any subset A of a topological space (X, τ^+) . The following statements are true.

- 1) $X \setminus B^+CCl^+(A) = B^+CInt(X \setminus A)$
- 2) $X \setminus B^+CInt(A) = B^+CCl^+(X \setminus A)$
- 3) $B^+CCl^+(A) = X \setminus B^+CInt(X \setminus A)$
- 4) $B^+CInt(A) = X \setminus B^+CCl^+(X \setminus A)$

Proof:

We only prove i), the other parts can be proved similarly.

For any point $x \in X$, $x \in X \setminus B^+CCl^+(A)$ implies that $x \notin B^+CCl^+(A)$.

Then for each $G \in B^+CO(X)$ containing x , $A \cap G = \emptyset$.

Then $x \in G \subset X \setminus A$, thus $x \in B^+CInt(X \setminus A)$.

References

- [1] D. Andrijevic, "On b-open sets", Mathmaticki. vesnik, 59-64, 48(1996).

- [2] H.Z.Ibrahim, "Bc-open sets in topological spaces", Advances in Pure Math., 3, 34-40, (2013).
- [3] F.Nirmalairudayam, "Bc-open sets in extended topological spaces", IJAR 2(9), 436-442, 2016.
- [4] A.S.Majid, "On some topological spaces by using b-open sets", M.S.C. Thesis University of AL-Qadissiya, college of mathematics and computer science, 2011.
- [5] N.Levine, "Semi-Open sets and Semi-Continuity in Topological Spaces," American Mathematical Monthly, Vol. 70. No. I. 1963, pp. 36-41. doi:10.2307/2312781.
- [6] N.V.Velicko. "H-closed Topological Spaces." American Mathematical Society. Vol. 38. No. 2, 1968, pp. 103-118.
- [7] R. H. Yunis, "Properties of θ -Semi-open sets." Zanco Journal Of Pure and applied sciences, Vol. 19, No. 1, 2007, pp. 116-122.
- [8] N.K. Ahmed, "On Some Types of Separation Axioms," M.Bc. Thesis, Salahaddin University, Arbil, 1990.
- [9] J. E. Joseph and M. H. Kwack, "On S-Closed Spaces," Bulletin of the American mathematical Society, Vol. 80, No. 2, 1980, pp. 341-348.
- [10] N.V. Velicko, "H-Closed Topological Spaces," American Mathematical Society, Vol. 78, No. 2, 1968, pp. 103-118
- [11] G. Di Maio and T. Noiri, "On s-Closed Spaces." Indian Journal of Pure and Applied Mathematics, Vol. 18, No. 3. 1987, pp. 226-233.
- [12] R. H. Yunis, "Properties of θ -Semi-open sets." Zanco Journal Of Pure and applied sciences, Vol. 19, No. 1, 2007, pp. 116-122.
- [13] M. H. stone, "Applications of the Theory of Boolean Rings to Topology," Transactions of the American mathematical Society, Vol. 41, No. 3, 1937, pp. 375-481. doi: 10.1090/S0002-9947-1937-1501905-7.
- [14] S. G. Crossley and S. K. Hildebrand, "semi-Closure," Texas Journal of Science, Vol. 22, No. 2-3, 1971, pp. 99-112.