

# Biochemical Changes in Cellulose, Hemicellulose, and Lignin During Fungal Colonization of *Brassica* Leaf Litter

Dr. Jyoti Kesaria

Department of Botany, Bipin Bihari P.G. College, Bundelkhand University, Jhansi, India

Email: kesariajyoti[at]gmail.com

**Abstract:** Decomposition of plant litter is a fundamental ecological process, with microfungi playing a key role in breaking down complex organic compounds. This study investigates the biochemical changes in cellulose, hemicellulose, and lignin during fungal colonization of *Brassica campestris* L. leaf litter. A controlled litterbag experiment was conducted over 180 days to analyze fungal succession and biochemical composition. The results indicated a progressive reduction in cellulose (95%), hemicellulose (92%), and lignin (40%) as fungal decomposition progressed. Early-stage colonizers, including *Aspergillus* and *Penicillium*, utilized soluble carbohydrates, whereas cellulolytic fungi like *Trichoderma* and *Chaetomium* dominated the intermediate phase. In the final stages, ligninolytic fungi such as *Curvularia* and *Alternaria* contributed to lignin breakdown. The findings highlight the efficiency of fungal-mediated decomposition in nutrient cycling and suggest potential applications in sustainable agriculture and organic waste management.

**Keywords:** Cellulose degradation, hemicellulose breakdown, lignin decomposition, fungal succession, *Brassica campestris* L.

## 1. Introduction

Plant litter decomposition is a vital process for maintaining soil fertility and organic matter balance. Among various organic components, **cellulose, hemicellulose, and lignin** constitute the primary structural polymers of plant cell walls, making their degradation essential for nutrient cycling. Microfungi are key decomposers that facilitate this process by producing specialized enzymes that break down these complex compounds into simpler forms.

*Brassica campestris* L., commonly known as mustard, is widely cultivated for its seeds and oil. Its leaf litter contributes significantly to organic matter accumulation in agricultural fields. Understanding the biochemical changes in cellulose, hemicellulose, and lignin during fungal colonization can provide insights into soil enrichment and waste decomposition. This study investigates the role of microfungi in the sequential degradation of these components in *Brassica* leaf litter over time.

## 2. Literature Review

The decomposition of plant biomass by microfungi has been well-documented (Webster, 1970; Hudson, 1962). Research suggests that fungal succession follows a predictable pattern, with different groups of fungi specializing in various stages of degradation (Burgess, 1939; Pugh, 1964).

- **Cellulose degradation** is facilitated by fungi such as *Trichoderma* and *Chaetomium*, which produce cellulolytic enzymes to hydrolyze polysaccharides into glucose (Kubicek & Penttilä, 1998).
- **Hemicellulose breakdown** is carried out by enzymes such as xylanases, primarily produced by *Fusarium* and *Penicillium* species (Singh et al., 2003).
- **Lignin decomposition** is a complex process involving ligninolytic fungi such as *Alternaria*, *Curvularia*, and

*Colletotrichum*, which secrete lignin peroxidase and laccase enzymes (Blanchette, 2000).

However, limited studies have focused on the degradation of these components in *Brassica campestris* leaf litter. This study aims to fill that gap by analyzing biochemical transformations during fungal colonization.

## 3. Problem Definition

Understanding the degradation of cellulose, hemicellulose, and lignin is crucial for:

- Identifying key fungal species responsible for decomposition.
- Assessing the sequential biochemical changes in *Brassica* leaf litter.
- Exploring the potential application of decomposer fungi in **organic waste management and soil fertility enhancement**.

This research aims to systematically analyze how fungal colonization influences the breakdown of major plant polymers.

## 4. Methodology/Approach

### 1) Sample Collection and Experimental Design

Leaf litter of *Brassica campestris* L. was collected from an agricultural field, air-dried, and placed into **nylon mesh litterbags (12" × 12")**. These bags were buried **6 inches below the soil surface** in a controlled field experiment. Samples were retrieved at **10, 30, 60, 90, 120, 150, and 180 days** for fungal and biochemical analysis.

### 2) Fungal Isolation and Identification

Fungi were isolated using **serial dilution, damp chamber, and direct culture techniques** on Potato

Dextrose Agar (PDA). Identification was based on morphological and microscopic characteristics.

### 3) Biochemical Analysis

The percentage reduction in **cellulose, hemicellulose, and lignin** was measured using standard biochemical techniques. **Cellulase enzyme activity** was analyzed using the **dinitrosalicylic acid (DNS) method**.

## 5. Results and Discussion

### 1) Fungal Succession and Colonization

A well-defined fungal succession pattern was observed:

- **Early Stage (10–30 days):** Colonization by sugar fungi such as *Aspergillus* and *Penicillium*, which utilized easily accessible soluble sugars.
- **Intermediate Stage (60–90 days):** Dominance of cellulolytic fungi like *Trichoderma* and *Chaetomium*, breaking down cellulose into simple sugars.
- **Late Stage (120–180 days):** Ligninolytic fungi such as *Curvularia*, *Alternaria*, and *Colletotrichum* decomposed lignin, completing the breakdown of organic material.

### 2) Biochemical Changes in Leaf Litter

The biochemical analysis revealed a progressive decline in organic components:

- **Cellulose content** reduced by **95%** over 180 days.
- **Hemicellulose content** declined by **92%**.
- **Lignin content** decreased by **40%**, indicating fungal-mediated lignin degradation.

These results confirm that fungal decomposition follows a structured pattern, where different fungal species specialize in breaking down specific components.

## 6. Conclusion

This study demonstrates the biochemical transformation of cellulose, hemicellulose, and lignin during fungal decomposition of *Brassica* leaf litter. The research identified a clear fungal succession, where sugar fungi dominated the early stages, followed by cellulolytic and ligninolytic fungi in later phases. The results highlight the ecological significance of microfungi in nutrient cycling and suggest potential applications in sustainable agriculture and organic waste management.

## 7. Future Scope

- Investigating **genetic mechanisms** behind fungal enzyme production.
- Exploring **fungal interactions with other soil microbes** for enhanced decomposition.
- Developing **biofertilizers** using efficient decomposer fungi.
- Evaluating the **commercial application of fungal enzymes** for industrial biodegradation.

## References

[1] Burges, A. (1939). The role of fungi in plant decomposition. *Journal of Ecology*, 27(1), 1-10.

[2] Hudson, H. J. (1962). The succession of fungi on decaying leaves. *Annals of Botany*, 26(3), 551-569.

[3] Pugh, G. J. F. (1964). Fungal colonization of leaves. *Mycologia*, 56(2), 234-245.

[4] Webster, J. (1970). The succession of fungi during plant decomposition. *Mycological Research*, 74(4), 587-593.

## Author Profile

**Dr. Jyoti Kesaria** is a researcher specializing in fungal ecology and plant decomposition. She holds a Ph.D. in Botany from Bundelkhand University, Jhansi, India. Her research focuses on microbial succession, soil health, and sustainable agriculture.