

# Haematological Response of *Rastrineobola argentea* Exposed To Subchronic Doses of Inorganic Cadmium (Cd), Methylmercury and Polychlorinated Biphenyl (PCB) via Dietary Exposure Pathway

Emily Jepyegon Chemoiwa<sup>1</sup>, Elijah Oyoo-Okoth<sup>2\*</sup>

<sup>1</sup>Department of Biological Sciences, University of Eldoret, P.O. Box 1125-30100, Eldoret, Kenya

<sup>2</sup>School of Natural Resources and Environmental Studies, Department of Natural Resource, Karatina University, P.O. Box 1957-10101, Karatina, Kenya

**Abstract:** Haematological indices provide information on various aspects of fish health when exposed to environmental contaminants. We evaluated the effects of cadmium, methyl mercury (MeHg), and Polychlorinated Biphenyl (PCB) on the haematological parameters of a tropical fish. One hundred and eighty (180) mature individuals of the freshwater fish *Rastrineobola argentea* were exposed to dietary subchronic doses of Cd ( $4 \mu\text{g g}^{-1} \text{CaCl}_2$ ), MeHg ( $0.1 \text{CH}_3\text{HgCl} \mu\text{g g}^{-1}$ ) and PCBs ( $0.2 \mu\text{g g}^{-1}$ ). After 45 days, blood was sampled from exposed and control groups to evaluate haematological effects of contaminants on erythrocytes, total leukocytes and differential leukocytes counts, haematocrit, haemoglobin concentration, red blood cell indices, mean corpuscular volume (MCV), mean corpuscular haemoglobin (MCH), and mean corpuscular haemoglobin concentration (MCHC). Results showed that red blood cells counts, haemoglobin concentration, haematocrit, leukocytes, neutrophils, and mononuclear cells counts, significantly ( $P < 0.05$ ) decrease between control groups compared with the Cd, MeHg and PCB tested groups. The MCV, MCH and MHC showed significant ( $P > 0.05$ ) increase between control groups with the Cd, MeHg and PCB tested groups. The present study shows that changes in haematological parameters were detectable at sub-chronic exposure to contaminants, but their application in field biomonitoring using *R. argentea* will need more detailed studies.

**Keywords:** Biomarkers; MeHg, PCB, Haematology; Tropical fish; *Rastrineobola argentea*

## 1. Introduction

The release of industrial, domestic, and urban wastes generated through anthropogenic activities into aquatic ecosystems normally cause stress to the aquatic life. These include metals and their methylated forms and organic contaminants. Currently, the widespread uses of metals, the legacies of past contamination and new technologies, continue to increase the concentration of metal into the aquatic environment [1]. Cadmium has no known roles in the fish and will be detrimental even in low exposure doses, is considered an ubiquitous toxicant and poses significant health risk in many parts of the world. This metal is one of the most commonly used metals in industry and its toxicity is of concern to public health due to its persistence in the environment [2,3]. Industrial uses of Cd and agricultural uses of phosphate fertilizers have caused widespread dispersion of the metal at trace levels into the environment and human foodstuffs [4,5]. Methyl mercury (MeHg) is a highly lipophilic environmental contaminant which easily crosses the blood barrier, and the primary route of exposure is through ingestion of contaminated food [6]. Polychlorinated biphenyls (PCBs) were first manufactured commercially in 1929 and used widely as electric insulators in transformers, hydraulic fluids and paint additives [7]. Serious concerns about the distribution of PCBs were raised since they were found to be ubiquitous and persistent in the environment and biota samples such as soil, water, animal and human tissues [8].

Although the production of PCBs has been banned since the early 1970s [9], PCBs persist as legacy pollutants in which the chronic toxicity still represents a serious environmental risk. These contaminants have different modes of action on the aquatic organism by exuding stress response in aquatic organisms. Thus, the assessment of environmental disturbances requires the elucidation of stress effects throughout the hierarchy of biological organization.

The use of haematological endpoints is reasonable biomarkers of fish health [10,11]. Knowledge of the haematological characteristics is an important tool that can be used as an effective and sensitive index to monitor physiological and pathological changes in fishes [12]. Normal ranges for various blood parameters in fish have been established by different investigators in fish physiology and pathology [13,14]. In addition, haematological studies provide quite frequently and routinely accepted procedures in fish diagnosis to evaluate the interactions between dietary levels of nutrients [15]. Although fish blood parameters have been increasingly determined in environmental monitoring programs as valuable indicators of physiological changes in the presence of toxicants, the most important barrier to using these findings in environmental studies is the lack of basic information about the blood response to stressors for many tropical species [16]. The aim of this study was to determine the effects of subchronic dietary exposures to

metals (Cd), its methylated form (MeHg) and organic contaminant (PCB) in a tropical fish *Rastrineobola argentea* by analysis of haematological biomarkers.

## 2. Materials and Methods

The experiments were approved by the animal welfare committee before starting the experiments. A total of about 180 *R. argentea* (mean weight =  $1.50 \pm 0.42$  g) were collected between April and July 2010 off the coast of Lake Victoria, Kenya ( $0^{\circ}12'40''S$  and  $34^{\circ}49'30''E$ ) and transported to the Kenya Marine and Fisheries Research (KEMFRI) Laboratory Kisumu in Kenya and acclimated to the experimental condition for 28 days (mean weight:  $23.50 \pm 0.42$  g). Fish were reared in race-way type water tanks supplied with filtered lake water (0.45- $\mu$ m filtered water). The renewal rate was  $24 \text{ L h}^{-1}$ ; salinity: 0.5‰; pH:  $7.4 \pm 0.4$ ; dissolved oxygen:  $> 5.0 \text{ mg l}^{-1}$ . Fish were fed commercial feed of protein level 32%.

After 28 days, feeds were laced with Cd ( $4 \mu\text{g g}^{-1} \text{ CaCl}_2$ ), MeHg ( $0.1 \mu\text{g g}^{-1} \text{ CH}_3\text{HgCl}$ , Sigma), and 2, 2', 4, 4', 5, 5' (PCB 153) ( $0.2 \mu\text{g g}^{-1}$ ) before the feeding experiments. The feeding was done for 45 days for a period lasting for 30 min in the morning (0800 h) and evening (1700 h). 20 individuals were kept as control groups.

After 45 days of feeding exposure, the individuals from each group (tested and control) were anesthetized with 0.02% MS222 (ethyl-ester.3.aminobenzoic acid, Sigma). Blood samples (1.5 mL) were taken by caudal puncture with heparinized syringes containing 0.1 mL of anticoagulant (after filling up and expelling about 1.0 mL), and no additional heparin was added to the Eppendorf tubes.

Numbers of erythrocytes (red blood cells (RBC) count,  $10^6$  cells  $\text{ml}^{-1}$ ) and leukocytes (Lk count  $10^4$  cells  $\text{ml}^{-1}$ ) were determined by the haemocytometer; haematocrit (Ht v/v ratio or %) was determined by the microhaematocrit method; haemoglobin concentrations (Hb g  $\text{dl}^{-1}$ ) were determined measurement by the cyanometahaemoglobin method [17]; and the leukocyte differential count was made in peripheral blood smears stained by Giemsa [18], giving the Neutrophils value of differential neutrophils  $\times (100 \text{ leukocytes count})^{-1}$  and the Mononuclear value of differential lymphocytes plus monocytes  $\times (100 \text{ leukocytes count})^{-1}$ . The total number of thrombocytes was estimated in relation to the numbers of red blood cells and the proportion of each cell type observed in the blood smears [19]. The red cell indices, mean corpuscular volume (MCV), mean corpuscular haemoglobin (MCH), and mean corpuscular haemoglobin concentration (MCHC) were calculated from RBC, Ht, and Hb using protocols of Lee et al. [17].

## 3. Results

The means of the haematological values with their respective standard errors ( $\pm$  SE) are presented in Table 1. Haematological parameters including: red blood cells counts, haemoglobin concentration, haematocrit, leukocytes, neutrophils, and mononuclear cells counts, showed significant ( $P < 0.05$ ) decrease between the control groups when compared with the Cd, MeHg and PCB tested groups (Table 1). Fish exposed to these contaminants showed reduced haematological parameters with most of the reduced blood parameters being recorded in fish exposed to MeHg. On the contrary, MCV, MCH and MHC showed significant ( $P > 0.05$ ) increase between control groups with the Cd, MeHg and PCB tested groups.

**Table 1:** Blood parameters for *Rastrineobola argentea* exposure to inorganic cadmium [ $4 \mu\text{g g}^{-1} \text{CaCl}_2$ ], methyl mercury [ $0.1 \text{ MeHg } \mu\text{g g}^{-1} \text{CH}_3\text{HgCl}$ ], and Polychlorinated biphenyls [ $0.1 \text{ PCB } \mu\text{g g}^{-1}$ ] for 45 days

	Control	Cd	MeHg	(PCB 153)
RBC ( $\times 10^6 \mu\text{l}$ )	$3.47 \pm 0.62$	$2.54 \pm 0.43$	$2.42 \pm 0.51$	$2.83 \pm 0.52$
Leukocyte counts ( $\times 10^4 \text{ cells ml}^{-1}$ )	$4.22 \pm 0.75$	$3.45 \pm 0.89$	$2.98 \pm 0.87$	$3.12 \pm 0.67$
Haemoglobin (%)	$7.23 \pm 0.64$	$6.02 \pm 0.94$	$5.32 \pm 0.89$	$5.41 \pm 0.55$
Haematocrit (%)	$33.22 \pm 2.88$	$27.88 \pm 1.11$	$26.03 \pm 1.01$	$26.33 \pm 0.97$
Neutrophils counts (100 leukocytes count) <sup>-1</sup>	$45.94 \pm 2.88$	$33.23 \pm 4.11$	$24.52 \pm 3.77$	$26.78 \pm 2.33$
Mononuclear cell counts (100 leukocytes count) <sup>-1</sup>	$73.23 \pm 6.22$	$60.12 \pm 4.55$	$59.89 \pm 4.66$	$64.78 \pm 3.89$
MCV ( $\text{mm}^3 \text{ cell}^{-1}$ )	$123.42 \pm 5.99$	$154.34 \pm 8.78$	$172.22 \pm 7.98$	$165.33 \pm 10.54$
MCH ( $\text{pg cell}^{-1}$ )	$17.12 \pm 2.44$	$21.33 \pm 4.22$	$25.33 \pm 5.92$	$24.55 \pm 3.32$
MCHC ( $\text{g dl}^{-1}$ )	$13.89 \pm 3.11$	$18.21 \pm 3.88$	$18.89 \pm 4.11$	$18.44 \pm 1.11$

MCV - Mean Corpuscular Volume; MCH - Mean Corpuscular Haemoglobin; MCHC - Mean Corpuscular Haemoglobin Concentration

#### 4. Discussion

There is consistent lack of information related to haematological analysis of fish chronically exposed to food contaminated by metals, and organic substances. Therefore the aim of the current study was to determine the sub-chronic dietary exposures to metals (Cd), its methylated form (MeHg) and organic contaminant (PCB) in a tropical fish *R. argentea* by analysis of haematological biomarkers. We recorded no mortality of fish during experiment suggesting that the levels used were indeed low to cause toxic effects in fish. The reduction in RBC and leucocyte counts suggests a reduction in the blood  $\text{O}_2^-$  carrying capacity to the pollutant pointing to the cytotoxic effects of the pollutants as reported for *Onchorhynchus mykiss* after an acute exposure to aluminum [20] and in *Labeo rohita* exposed to sublethal levels of cypermethrin and carbofuran [12]. Nevertheless, other mechanisms of toxicity may be associated with the  $\text{O}_2^-$  carrying capacity such as the inhibition of iron absorption and defective iron metabolism shortening the life span in erythrocytes [21]. The reduced leukocyte counts could also be related to the presence of tecidual damages such as necrosis [22]. The values observed for haematocrit and Hb in *R. argentea* from control groups are relatively close to those of other tropical and nontropical species of fish such as *Ictalurus punctatus* (23.9%) [15] and *Colossoma macropomum* (20–23%). After exposure to pollutants, the haematocrit values of fish have been reported to reduce [10, 23]. Therefore the large reduction in haematocrit values when fish was exposed to MeHg after trophic subchronic exposure, indicate that MeHg may be more toxic to the cells. These results do not agree with those of Chowdhury et al. [24] and Oliveira Ribeiro et al. [25] who noted an increase of blood haematocrit and haemoglobin during environmental hypoxia and chronic or acute exposure to waterborne metals (Cd, Zn, Cu, Al, and Ni) to increase blood oxygen carrying capacity when impairment of gas exchange occurs. It is known that changes in leukocyte counts after exposure to pollutants may be associated to a decrease in

nonspecific immunity of the fish. In the current work the effects of MeHg on the leukocytes counts were among the most evident compared with those of other tested metals and control groups.

The increased MCV, MCH and MCHC observed in individuals of *R. argentea* exposed to MeHg may be explained by the presence of a larger amount of older or larger red blood cells as described by Hardig and Hoglund [26]. In addition, the exposure to mainly MeHg could also affect the mechanism of red blood cell turnover, increasing the number of circulating older cells and inducing an anemic state [12]. Some studies have pointed out a chronic effect of MeHg in head kidney of *Hoplias malabaricus*, indicating severe damage to this haematopoietic tissue [25] and supporting the hypothesis of a failure in the red blood cells turnover.

#### 5. Conclusion

Haematological parameters could be useful to evaluate the effects of contaminants. The present results showed that under experimental conditions blood parameters were sensitive to different aspects of contaminant exposure. Also, its ability to adapt to experimental conditions, its voracious behavior and its food chain position make *R. argentea* an interesting model to be used in experiments testing dietary exposure to contaminants but the application of these findings to preparation of environmental diagnoses will need a more investigation and must be validated in situ before establishing them as biomarkers.

#### References

- [1] Luoma, S.M., Rainbow, P.S., 2008. Metal contamination in aquatic environments: Science and lateral management. Cambridge University Press: Cambridge. 573pp.

- [2] Ogunseitan, O.A., Yang, S., Ericson, J., 2000. Microbial d-aminolevulinic dehydratase as a biosensor of lead bioavailability in contaminated environments. *Soil Biol. Biochem.*, 32, 1899–1906.
- [3] Gurer, O.H., Sabýy, H.U., Özgünes, H., 2004. Correlation between clinical indicators of lead poisoning and oxidative stress parameters in controls and lead-exposed workers. *Toxicology*, 195, 147–154.
- [4] Satarug, S., Baker, J.R., Urbenjapol, S., et al., (2003). A global perspective on cadmium pollution and toxicity in non-occupationally exposed population. *Toxicol. Lett.*, 137, 65–83.
- [5] WHO/IPCS (1992). Cadmium. *Environmental Health Criteria Document 134*, pp. 1–280.
- [6] Limke, T.L., Heidemann, S.R., Atchison, W.D., 2004. Disruption of intraneuronal divalent action regulation by methylmercury: are specific targets involved in altered neuronal development and cytotoxicity in methylmercury poisoning? *Neurotoxicology*, 25, 741–760.
- [7] Safe, S.H., 1994. Polychlorinated biphenyls (PCBs): environmental impact, biochemical and toxic responses, and implications for risk assessment. *Crit. Rev. Toxicol.*, 24, 87–149.
- [8] Jansson, B., Andersson, R., Asplund, L., et al., 1993. Chlorinated and brominated persistent organic compounds in biological samples from the environment. *Environ. Toxicol. Chem.*, 12, 1163–1174.
- [9] Harrad, S.J., Sewart, A.P., Alcock, R., et al., 1994. Polychlorinated biphenyls (PCBs) in the British environment: sinks, sources and temporal trends. *Environ. Pollut.* 85, 131–146.
- [10] Lohner, T.W., Reash, R.J., Willet, V.E., et al., 2001. Assessment of tolerant sunfish populations (*Lepomis* sp.) inhabiting selenium-laden coal ash effluents. *Ecotoxicol. Environ. Saf.*, 50, 203–216.
- [11] Satheeshkumar, P., Ananthan, G., Senthil Kumar, D., et al. 2011. Haematology and biochemical parameters of different feeding behaviour of teleost fishes from Vellar estuary, India. *Comp Clin Pathol.* DOI 10.1007/s00580-011-1259-7.
- [12] Adhikari, S., Sarkar, B., Chatterjee, A., Mahapatra, C.T., et al., 2004. Effects of cypermethrin and carbofuran on certain haematological parameters and prediction of their recovery in a freshwater teleost; *Labeo rohita* (Hamilton). *Ecotoxicol. Environ. Saf.*, 58, 220–226.
- [13] Rambhaskar B, Srinivasa Rao K (1986) Comparative haematology of ten species of marine fish from Visakhapatnam Coast. *J. Fish Biol.* 30, 59–66
- [14] Xiaoyun, Z., Mingyun, L., Khalid, A., et al., 2009. Comparative of haematology and serum biochemistry of cultured and wild Dojo loach *Misgurnus anguillicadatus*. *Fish Physiol. Biochem.*, 35:435–441.
- [15] Lim, C., Klesius, P.H., Li, M.H., Robinson, E.H., 2000. Interaction between dietary levels of iron and vitamin C on growth, haematology, immune response and resistance of channel catfish (*Ictalurus punctatus*) to *Edwardsiella ictaluri* challenge. *Aquaculture*, 185, 313–327.
- [16] Affonso, E.G., Polez, V.L.P., Corrêa, C.F., et al., 2002. Blood parameters and metabolites in the teleosts fish *Colossoma macropomum* exposed to sulfide or hypoxia. *Comp. Biochem. Physiol. C*, 133, 375–382.
- [17] Lee, R.G., Foerster, J., Jukens, J., et al., 1998. *Wintrobe's—Clinical Haematology*, 10<sup>th</sup> ed. Lippincott Williams & Wilkins, New York, USA.
- [18] Beutler, E., Lichtman, M.A., Coller, B.S., et al., 2001. *Hematology*, sixth ed. McGraw-Hill, USA.
- [19] Carvalho, W.F., 1994. *Técnicas médicas de hematologia e imunohaematologia*. Editora Coopmed, Belo Horizonte, Brasil, pp. 66–175.
- [20] Allin, C.J., Wilson, R.W., 2000. Effects of pre-acclimation to aluminium on the physiology and swimming behaviour of juvenile rainbow trout (*Oncorhynchus mykiss*) during a pulsed exposure. *Aquat. Toxicol.*, 51, 213–224.
- [21] Liu, J., Liu, Y., Habeebu, S.S., et al., 1999. Metallothionein-null mice are highly susceptible to the haematotoxic and immunotoxic effects of chronic  $\text{DcCl}_2$  exposure. *Toxicol. Appl. Pharmacol.*, 159, 98–108.
- [22] Oliveira Ribeiro, C.A., Belger, L., Pelletier, E., et al., 2002. Histopathological evidence of inorganic mercury and methylmercury toxicity in the arctic charr (*Salvelinus alpinus*). *Environ. Res.*, 90, 217–225.
- [23] Mattsson, K., Lehtinen, D.-J., Tana, J., et al., 2001. Effects of pulp mill effluents and restricted diet on growth and physiology rainbow trout (*Oncorhynchus mykiss*). *Ecotoxicol. Environ. Saf.*, 49, 144–154.
- [24] Chowdhury, M.J., McDonald, D.G., Wood, C.C., 2004. Gastrointestinal uptake and fate of cadmium in rainbow trout acclimated to sublethal dietary cadmium. *Aquat. Toxicol.*, 69, 149–163.
- [25] Oliveira Ribeiro, C.A., Filipak Neto, F., Mela, M., et al., 2006. Haematological findings in neotropical fish *Hoplias malabaricus* exposed to subchronic and dietary doses of methylmercury, inorganic lead, and tributyltin chloride. *Environ. Res.*, 101, 74–80.
- [26] Hardig, J., Hoglund, L.B., 1983. Seasonal and ontogenetic effects on methaemoglobin and reduced glutathione content in the blood of reared Baltic salmon. *Comp. Biochem. Physiol.*, 75, 27–34

### Author Profile



**Dr. Emily Jepyegon Chemoiwa** has a PhD in Biological Science from University of Eldoret, Kenya and University of Kwazulu Natal, South Africa. She has diverse knowledge in Genetics, Evolutionary Biology, Limnology, Fish Population Dynamics and Toxicology.



**Dr. Elijah Oyoo-Okoth** has a PhD in Aquatic Ecology and Ecotoxicology from the University of Amsterdam, The Netherlands. He has published several publications in top ranking Toxicology Journals. He has diverse interested in Environmental Biology, Environmental Health, Fisheries Science and Ecotoxicology.