

# Synthesis, Anticancer and Anti-Microbial Characterization of Some Novel Quinazoline Derivatives

Manish Chaudhari<sup>1</sup>, J.J.Vora<sup>2</sup>

<sup>1</sup>Department of Chemistry, Mehsana Urban Institute Of Sciences , Ganpat University, Kherva

<sup>2</sup>Department of Chemistry, Hemchandracharya North Gujarat University, Patan

**Abstract:** With an objective to synthesize compounds with high anti-cancer and other anti-microbial properties, a systematically planned organic synthesis was carried out. Quinazoline has been a very significant heterocycle and also folic acid is having highly significant physiological activity. It was thought that combination of these two fragments must give resistant molecules with highly enhanced biological properties. With this ambiguous objective, the organic synthesis was carried out. The synthesized compounds were characterized by sensitive instrumental methods like Mass Spectra, <sup>13</sup>C NMR, Infra-Red Spectra, UV Spectra etc. Their structures were thus confirmed by different physicochemical methods.

**Keywords:** Anticancer Activity, Antimicrobial Activity, Quinazoline derivatives and Amino Acid Derivatives

## 1. Introduction

Quinazoline and their derivatives constitute an important class of heterocyclic compounds. Many of them show insecticidal, analgesic, antifungal, antibacterial, anticancer, anti-inflammatory activities. Quinazoline nucleus is found in many bioactive natural products. Because of these reasons much attention is being paid for the synthesis of quinazoline derivatives. Looking at the biological significance of quinazoline nucleus it was thought to design and synthesize new quinazoline derivatives and screen them for their antibacterial activity.


## 2. Materials and Method

Synthesis of quinazoline derivatives involved following steps: In the first step anthranilic acid was treated with benzoyl chloride to give 2-phenyl-3,1 benzoxazine - 4 [3H] one. In the next step 2-phenyl-3,1 benzoxazine - 4 [3H]


one was allowed to react with hydrazine hydrate to give 3-amino-2-phenyl quinazoline-4(3H)-one which was further treated with methionine and proline by the method given.

### General procedure for synthesis of various 3- amino 2-phenyl quinazoline 4(3H) one and amino acid derivative.

3-amino 2-phenyl quinazoline 4(3H) one (0.0253M) in different amino acid (0.0253M) are taken and mixed to give a homogeneous mixture. This reaction mixture was taken in petri dish and it was placed in microwave oven and full microwave was set and heated for 5 min. in grilled position and after that the product was collected.




Scheme: 2



3-amino 2-phenyl quinazoline 4(3H) one and Methionine derivative

Scheme: 3



3-amino 2-phenyl quinazoline 4(3H) one and Proline derivative

### 3. Spectra Characterization

**Compound-1:** 3-amino 2-phenyl quinazoline 4(3H) one and Methionine derivative

**Elemental Analysis:** %C %H %N %S

**Found** 59.05 5.91 13.96 8.78

**Calculated** 57.51 5.85 14.02 12.82

**Infrared spectral Features (cm<sup>-1</sup>): cm<sup>-1</sup> Assignment**

1637 C=N Stretching  
 3213,3307 -CONH<sub>2</sub>, -NH- Stretching  
 1563, 1588, 1611 -CONH<sub>2</sub>, -NH- Bending  
 1661 C=O Stretching  
 2915 Aromatic C-H Stretching  
 1588, 1611 C=C Stretching

1448, 1370 Alkenes -CH<sub>3</sub> bending

1472 Alkenes -CH<sub>2</sub>- bending

2577 S-H, -S- Stretching

**<sup>13</sup>C Spectral Features (ppm): (ppm) Assignment**

38.55, 39.27, 39.48, 39.69, 39.90, 40.10 R<sub>2</sub>-CH<sub>2</sub>, R<sub>3</sub>-CH, R<sub>4</sub>-C

78.50, 78.83, 79.16 C-N

146.67, 155.50 R-CO-NH

125.95, 126.50, 126.88 127.30, 127.37, 129.52,

134.03, 134.76 Benzene

161.11, 167.80 =C=O

**Mass Spectral Features: (ppm) Assignment**

499 Molecular peak is observable

500 (M+1) Molecular peak is observable

501 (M+2) molecular peak is observable

483 Base peak is observable for (M - CH<sub>2</sub>)

278 Peak is observable for (M – C<sub>8</sub>H<sub>17</sub>N<sub>2</sub>S<sub>2</sub>O)  
 238 Peak is observable for (M – C<sub>10</sub>H<sub>17</sub>N<sub>2</sub>S<sub>2</sub>O<sub>2</sub>)  
 260 Peak is observable for (M - C<sub>8</sub>H<sub>17</sub>N<sub>2</sub>S<sub>2</sub>O.H<sub>2</sub>O)  
 150 Peak is observable for Methionine (M+1)

**UV Spectral Features:  $\lambda_{\max}$  Assignment**

289 n $\rightarrow$  $\pi^*$  Substituent with lone pair R-band  
 248  $\pi\rightarrow\pi^*$  Substituent Delocalized by Aryl K-band  
 215,221  $\pi\rightarrow\pi^*$  Allowed

**Compound – 2:** 3-amino 2-phenyl quinazoline 4(3H) one and Proline derivative

**Elemental Analysis :**

%C %H %N

**Found** 63.24 5.62 15.23

**Calculated** 66.82 6.08 16.24

**Infrared spectral Features (cm<sup>-1</sup>) :****cm<sup>-1</sup> Assignment**

1635 C=N Stretching  
 3214,3307, 3358, 3442, 3507 -CONH<sub>2</sub>, -NH- Stretching  
 1563, 1574, 1590, 1611 -CONH<sub>2</sub>, -NH- Bending  
 1661 C=O Stretching  
 3035, 3061 Aromatic C-H Stretching  
 1590, 1611 C=C Stretching

1448, 1371 Alkenes –CH<sub>3</sub> bending

1471 Alkenes –CH<sub>2</sub>- bending

**<sup>13</sup>C Spectral Features (ppm) :****(ppm) Assignment**

22.83, 2379, 27.16, 28.90 R-CH<sub>3</sub>  
 38.89,39.31,39.73,40.14, 44.45, 45.15, 59.73,60.49 R<sub>2</sub>-CH<sub>2</sub>,  
 R<sub>3</sub>-CH, R<sub>4</sub>-C  
 78.48,78.81, 79.14 C-N  
 146.67,155.44 R-CO-NH  
 125.94, 126.44, 127.27,128.49,129.34, 131.15,  
 131.35, 133.97, 134.75 Benzene  
 161.08, 164.23, 165.87 =C=O

**Mass Spectral Features :****(ppm) Assignment**

335 (M+1) Molecular peak is observable  
 238 Peak is observable for (M – C<sub>5</sub>H<sub>6</sub>NO) Quinazoline peak  
 116 Peak is observable for Proline (M+1)

**UV Spectral Features :** **$\lambda_{\max}$  Assignment**

290 n $\rightarrow$  $\pi^*$  Substituent with lone pair R-band  
 252  $\pi\rightarrow\pi^*$  Substituent Delocalized by Aryl K-band  
 206,220  $\pi\rightarrow\pi^*$  Allowed

**Table 1:** Various 3-amino 2-phenyl quinazoline (3H) one derivative

Sr. No.	Compound Name	M.P	Solubility	Nitrogen Rule Obeyed	Rules of 13		Compound Formula	Base formula	Unsaturation Index U
					n	r			
1	Comp. 1	246	DMSO	yes	28	4	C <sub>24</sub> H <sub>29</sub> N <sub>5</sub> O <sub>3</sub> S <sub>2</sub>	C <sub>38</sub> H <sub>43</sub>	23
2	Comp. 2	176	DMSO	yes	25	9	C <sub>19</sub> H <sub>18</sub> N <sub>4</sub> O <sub>2</sub>	C <sub>25</sub> H <sub>34</sub>	13

**4. Antimicrobial Activity**

Antibacterial activity was performed by cup plate method by measuring zone of inhibition. All the test compounds were screened for antibacterial activity against bacterial strains *Staphylococcus aureus*, *Bacillus sp.*, *Salmonella typhi* and *Escherichia coli* (ESS 2231) at a concentration of 200, 300, 400  $\mu$ g/ml. Ampicillin was used as standard drug at a concentration of 200, 300, 400  $\mu$ g/ml, Nutrient agar was used as culture medium & DMF was used as solvent control.

The plates were inoculated within minutes of the preparation of suspension, so that the density does not change. A sterile cotton swab over was dipped into the suspension and the medium was inoculated by even streaking of the swab over the entire surface of the plate in three directions. After the inoculums had dried, cups of diameter 6mm were made in the agar plate with a sterile cork borer. The drugs solutions were added to these cups with a micropipette and the plates were then incubated at 37 °C for 24 hours. The zone of inhibition was measured using mm scale.

The fungicidal activity of all the compounds was studied at 1000ppm concentration. In vitro plant pathogenic organisms used were *Aspergillus* SP, Yeast, etc with standard antifungal, antibiotic fluconazol. The antifungal activity of all the compounds was measured on each of these plant

pathogenic strains on a potato dextrose agar (PDA) medium such a PDA medium contained potato 200gm, dextrose 20gm, agar 20gm, and water 1 liter. Five days old cultures were employed. The compounds to be tested were suspended (1000ppm) in a PDA medium and autoclaved at 120°C for 15 min. and at 15 atm pressure. These media were poured into sterile Petri plates and the organisms were inoculated after cooling the Petri plates. The percentage inhibition for fungi was calculated after five days using the formula given below.

$$\text{Percentage of Inhibition} = 100(X-Y)/X$$

Where, X = Area of colony in control plate Y= Area of colony in test plate

**Table 2:** Antibacterial Activity of Compounds

Compound Name	Zone of Inhibition (in mm)											
	Gram Positive			Gram Negative								
	Bacillus Sp.			Staphylococcus Aureus			E. Coli			Salmonella Typhi		
	200 $\mu$ g/ml	300 $\mu$ g/ml	400 $\mu$ g/ml	200 $\mu$ g/ml	300 $\mu$ g/ml	400 $\mu$ g/ml	200 $\mu$ g/ml	300 $\mu$ g/ml	400 $\mu$ g/ml	200 $\mu$ g/ml	300 $\mu$ g/ml	400 $\mu$ g/ml
Comp-1	18	23	22	25	28	30	22	25	28	15	19	24
Comp-2	17	23	28	--	--	18	--	--	17	--	18	24

**Table 3:** Antifungal Activity of Compounds

Compound Name	Zone of Inhibition ( in mm)					
	Aspergillus Sp.			Yeast		
	200 µg/ml	300 µg/ml	400 µg/ml	200 µg/ml	300 µg/ml	400 µg/ml
Compound- 1	13	18	23	18	22	25
Compound- 2	--	18	22	24	27	30

## 5. Anti Cancer Experimental Setup

### 5.1 Cell Lines and Culture Medium

MDA-MB-468 and HCT-15 cultures used in these experiments were derived from National Centre for Cell Science (NCCS), Pune. Stock cells of these cell lines were cultured in DMEM, supplemented with 10% FBS (fetal bovine serum). Along with media cells were also supplemented with 5% HBSS, penicillin, streptomycin and Amphotericin – B, in a humidified atmosphere of 5 % CO<sub>2</sub> at 37°C until confluence reached. The cells were dissociated with 0.2% trypsin, 0.02% EDTA in phosphate buffer saline solution. The stock cultures were grown initially in 25 cm<sup>2</sup> tissue culture flasks, than in 75 cm<sup>2</sup> and finally in 150 cm<sup>2</sup> tissue culture flask and all cytotoxicity experiments were carried out in 96 microtitre well- plates. 2×10<sup>4</sup> cells/well was added in to each well of 96 well-plates.

**Table 4:** Results of MDA-MB-468 cell line

Conc. µM/ml	Log con.	% Cell Inhibition		
		Compound - 1	Compound - 2	Std. Doxorubicin
0.01	-2.29	-2.20	-5.06	-40.87
0.02	-1.82	-1.43	-7.01	-32.81
0.05	-1.34	-1.72	-7.01	-29.69
0.14	-0.86	-0.34	0.26	-17.63
0.41	-0.39	10.88	0.95	-14.34
1.23	0.09	13.17	0.84	-12.55
3.70	0.57	22.68	2.65	11.25
11.11	1.05	28.09	6.97	48.35
33.33	1.52	39.14	10.22	71.32
100.00	2.00	57.18	35.32	86.69
IC <sub>50</sub> (µM/ml)		31.96	>100	6.802
R <sup>2</sup>		0.9184	0.9593	0.9833

## 6. Result and Discussions

Antibacterial activity of compound 1 is good at lower as well as higher concentrations against E. Coli, S. aureus and bacillus sp. whereas against S. typhi, A. sp. and yeast is moderate to low in comparison with standard antibiotic. Compound 2 shows good activity against B. sp. and yeast. For other organisms, it shows less activity compared to the standard antibiotic.

Potent compounds with good cytotoxicity but less than std. drug.>M15. While no cytotoxicity effect by M17 compounds because of Ic50 values is more than 100. Therefore it would be better to go for further study of the non-toxic compound series after checking of its toxicity with normal cell line and then one can go for establishing mechanism based study of that series of compounds by using Tunnel assay, Flow cytometry, DNA fragmentation assay Or CASPACASE assay

## References

- [1] Theivendren panner selvam, Palanirajan Vijayarajkumar A review article of quinazoline Marketed drugs. Research in Pharmacy 1(1): 1-21, (2011)
- [2] Vijaychand. A, S.N.Manjula, Bharath. E.N. and Divya B. Review article of Medicinal and Biological significance of quinazoline. International Journal of Pharma and BioSciences. Vol-2, Issue-1 (Jan-Mar-2011)
- [3] R.S.Varma Solvent-free organic syntheses. Using supported reagents and microwave irradiation Green Chemistry: 1; 43-55, 1999
- [4] R.N.Gedye, F.E.Smith, K.C.Westaway The rapid synthesis of organic compounds in microwave ovens CAN.Journal of Chemistry: 66; 17-26, 1988
- [5] W. Kemp Organic Spectroscopy, ELBS (1996)
- [6] B. Oteorge and P. Mc. Intyre Infrared Spectroscopy, Heyden, London (1972)
- [7] L. J. Bellamy The Infrared Spectra of complex molecules, Methuen, London (1980)
- [8] R. M. Silverstein, G. Clayton Bassler and Terence C. Morrill Spectrometric Identification of Organic Compounds, Fifth Edition, John Wiley & Sons (1991)
- [9] G. M. Lampman, Donald L. Pavia, G. S. Kriz and J. R. Vyvyan Spectroscopy 4e, Fourth Edition, CENGAGE Learning. (2010)
- [10] M Suggitt, MC Bibby - Clinical Cancer Research, 2005 50 years of preclinical anticancer drugs screening : Empirical to target- driven approaches Clinical Cancer Research, 11; 971, (2005)
- [11] KJ Bussey, K Chin, S Lababidi, M Reimers Integrating data on DNA copy number with gene expression levels and drug sensitivities in the NCI-60 cell line panel. Molecular Cancer Therapeutics; 5: 853-67. (2006)
- [12] B. Willium "The textbook of Microbiology" W.B. Saunders Co., London, 16th edition. pp. 12 and pp 145 (1945)
- [13] W. Robert and E.G. Scott "Diagnostic Microbiology" The C.V. Mosby Co., Saint Louis, 2<sup>nd</sup> edition, pp 318 (1966)